arXiv:1409.5006v1 [math.CO] 17 Sep 2014

Extraction of the i^{th} Elementary Symmetric Polynomial from a Product in Binomial Form $\binom{m_1+...+m_n}{i}$

Felix de la Fuente *

September 17, 2014

Abstract

The i^{th} elementary symmetric polynomial of the set of n variables $\mathcal{R} = \{m_1, m_2, m_3, \ldots, m_n\}$ is isolated from the expansion of the i^{th} binomial product $\binom{m_1+...+m_n}{i}$ via an alternating sum.

1 Introduction

For a finite root set $\mathcal{R} = \{m_1, m_2, m_3, \ldots, m_n\}$ of any positive numbers $m_i \in \mathbb{N}$, whether equal or unequal, the i^{th} **elementary symmetric polynomial** $e_i(n)$ is the sum of all i-tuples of unique elements from $\mathcal{R}$

$$\begin{aligned} e_i(n) &= \sum_{1 \leqslant j_1 < j_2 < \ldots < j_i \leqslant n} m_{j_1} m_{j_2} \ldots m_{j_{i-1}} m_{j_i} \\ &= \sum_{J \in \binom{1,2,\ldots n}{i}} \prod_{j \in J} m_j \end{aligned}$$

When the size of the tuples equals 0, the polynomial accounts for the empty product $e_0(n) = 1$.

In this way, the set of four numbers $\mathcal{R} = \{m_1, m_2, m_3, m_4\}$ generates the following polynomials

$e_0(4) = 1$
$e_1(4) = m_1 + m_2 + m_3 + m_4$
$e_2(4) = m_1m_2 + m_1m_3 + m_1m_4 + m_2m_3 + m_2m_4 + m_3m_4$
$e_3(4) = m_1m_2m_3 + m_1m_2m_4 + m_1m_3m_4 + m_2m_3m_4$
$e_4(4) = m_1m_2m_3m_4$

Algebraically, elementary symmetric polynomials constitute a basis of the space of symmetric polynomials, and so do others functions such as homogeneous symmetric polynomials and power sum symmetric polynomials, which will not be discussed here. Relations among bases is a classical subject dating back to Guirard, Waring, Newton and Euler [1] (see [2] for a general overview) as it is the specialization due to the insertion of particular numbers, or q-numbers, within the root set. Together with the homogeneous symmetric polynomials, many of these specializations have historically been objects of interest in number theory and combinatorics for diverse authors, and still are focus of current research. Some noteworthy examples are illustrated in Figure 1.1.

*mail@felixdelafuente.es



*indicates a specialization over a root vector with the same number of elements

Figure 1.1: Some specialized root sets generating orthogonal families of numbers as elementary and homogeneous symmetric polynomials.

Let $\binom{N}{i}=\frac{N(N-1)\cdots(N-i+1)}{i!}$ and $\left(\!\binom{N}{i}\!\right)=\frac{N(N+1)\cdots(N+i-1)}{i!}$ be **binomial numbers** of the **first** and the **second kind** respectively.

If we consider N as the accumulate of the root set $N=\sum_{j=1}^{n} m_j$, the former binomial numbers represent the i^{th} **binomial product** $\binom{m_1+\ldots+m_n}{i}=\frac{(m_1+\ldots+m_n)(m_1+\ldots+m_n-1)\cdots(m_1+\ldots+m_n-i+1)}{i!}$ for $n>i$.

This article presents an expression for the elementary symmetric polynomials of a root set $\mathcal{R}=\{m_1, m_2, m_3, \ldots, m_n\}$ as a function of its associated binomial products. Namely

$$
\begin{aligned}
e_1(n) &= \tbinom{m_1+\ldots+m_n}{1} \\
e_2(n) &= \tbinom{m_1+\ldots+m_n}{2} - \left(\!\tbinom{n-1}{0}\!\right)\left[\tbinom{m_1}{2}+\tbinom{m_2}{2}+\tbinom{m_3}{2}+\tbinom{m_4}{2}+\cdots+\tbinom{m_n}{2}\right] \\
e_3(n) &= \tbinom{m_1+\ldots+m_n}{3} - \left(\!\tbinom{n-2}{0}\!\right)\left[\tbinom{m_1+m_2}{3}+\tbinom{m_1+m_3}{3}+\tbinom{m_2+m_3}{3}+\cdots\right] \\
&\quad + \left(\!\tbinom{n-2}{1}\!\right)\left[\tbinom{m_1}{3}+\tbinom{m_2}{3}+\tbinom{m_3}{3}+\tbinom{m_4}{3}+\cdots+\tbinom{m_n}{3}\right] \\
e_4(n) &= \tbinom{m_1+\ldots+m_n}{4} - \left(\!\tbinom{n-3}{0}\!\right)\left[\tbinom{m_1+m_2+m_3}{4}+\tbinom{m_1+m_2+m_4}{4}+\tbinom{m_1+m_3+m_4}{4}+\cdots\right] \\
&\quad + \left(\!\tbinom{n-3}{1}\!\right)\left[\tbinom{m_1+m_2}{4}+\tbinom{m_1+m_3}{4}+\tbinom{m_2+m_3}{4}+\cdots\right] \\
&\quad - \left(\!\tbinom{n-3}{2}\!\right)\left[\tbinom{m_1}{4}+\tbinom{m_2}{4}+\tbinom{m_3}{4}+\tbinom{m_4}{4}+\cdots+\tbinom{m_n}{4}\right] \\
e_5(n) &= \tbinom{m_1+\ldots+m_n}{5} - \left(\!\tbinom{n-4}{0}\!\right)\left[\tbinom{m_1+m_2+m_3+m_4}{5}+\tbinom{m_1+m_2+m_3+m_5}{5}+\tbinom{m_1+m_3+m_4+m_5}{5}+\cdots\right] \\
&\quad + \left(\!\tbinom{n-4}{1}\!\right)\left[\tbinom{m_1+m_2+m_3}{5}+\tbinom{m_1+m_2+m_4}{5}+\tbinom{m_1+m_3+m_4}{5}+\cdots\right] \\
&\quad - \left(\!\tbinom{n-4}{2}\!\right)\left[\tbinom{m_1+m_2}{5}+\tbinom{m_1+m_3}{5}+\tbinom{m_2+m_3}{5}+\cdots\right] \\
&\quad + \left(\!\tbinom{n-4}{3}\!\right)\left[\tbinom{m_1}{5}+\tbinom{m_2}{5}+\tbinom{m_3}{5}+\tbinom{m_4}{5}+\cdots+\tbinom{m_n}{5}\right]
\end{aligned}
$$

2 Extraction of the i^{th} Elementary Symmetric Polynomial

Theorem. *Let $\mathcal{R}=\{m_1,m_2,m_3,\ldots,m_n\}$ be a root set of any positive numbers $m_j\in\mathbb{N}$. Then, the i^{th} elementary symmetric polynomial of size n is given by*

$$e_i(n)=\binom{\sum_{k=1}^{n}m_k}{i}-\sum_{h=1}^{i-1}(-1)^{h-1}\left(\binom{n-i+1}{h-1}\right)\left[\sum_{J\in\binom{1,2,..n}{i-h}}\binom{\sum_{j\in J}m_j}{i}\right]$$

Proof. Consider the root set of n elements $\mathcal{R}=\{m_1,m_2,m_3,\ldots,m_n\}$ and a subset of s elements $\mathcal{Q}_s=\{m_{j_1},m_{j_2},\ldots,m_{j_s}\}$, $\mathcal{Q}_s\in\mathcal{R}$. The expansion of the product in binomial form $\binom{m_1+\ldots+m_n}{i}=\frac{(m_1+\ldots+m_n)(m_1+\ldots+m_n-1)\cdots(m_1+\ldots+m_n-i+1)}{i!}$ can be organized by association to sets of sums involving products whose factors belong to certain subset $\mathcal{Q}_s$.

Example: Consider the expansion of the 4^{th} binomial product $\binom{m_1+\ldots+m_n}{4}$. The summands involving products of elements belonging to a general subset of size two $Q_2=\{m_{j_1},m_{j_2}\}$ are

$$\begin{aligned}&+\left[-\tfrac{22}{4!}m_{j_1}m_{j_2}\right]\\&+\left[\tfrac{18}{4!}\left(m_{j_1}\right)^2m_{j_2}+\tfrac{18}{4!}m_{j_1}\left(m_{j_2}\right)^2\right]\\&+\left[-\tfrac{4}{4!}\left(m_{j_1}\right)^3m_{j_2}-\tfrac{6}{4!}\left(m_{j_1}\right)^2\left(m_{j_2}\right)^2-\tfrac{4}{4!}m_{j_1}\left(m_{j_2}\right)^3\right]\end{aligned}$$

The sum of the exponents of the factors of every product add up to an accumulate power p which ranges from the size of the set s to the maximal value i. The repetitions of every product are given by the coefficient $K_p^{\lambda(p)_k}$, where $\lambda(p)_k$is the ordered partition of p into i parts corresponding to the distribution of the exponents. Extending the former example for the expansion of the general i^{th} binomial product $\binom{m_1+\ldots+m_n}{i}$, the summands involving factors from $\mathcal{Q}_2$ are

$$\begin{aligned}&+\left[K_2^{(1,1)}m_{j_1}m_{j_2}\right]\\&+\left[K_3^{(2,1)}\left(m_{j_1}\right)^2m_{j_2}+K_3^{(1,2)}m_{j_1}\left(m_{j_2}\right)^2\right]\\&+\left[K_4^{(3,1)}\left(m_{j_1}\right)^3m_{j_2}+K_4^{(2,2)}\left(m_{j_1}\right)^2\left(m_{j_2}\right)^2+K_4^{(1,3)}m_{j_1}\left(m_{j_2}\right)^3\right]\\&\vdots\\&+\left[\textstyle\sum_k K_p^{\lambda(p)_k}(m_{j_1}m_{j_2})_p^{\lambda(p)_k}\right]\\&\vdots\\&+\left[\textstyle\sum_k K_i^{\lambda(i)_k}(m_{j_1}m_{j_2})_i^{\lambda(i)_k}\right]\\&=\textstyle\sum_{p=2}^{i}\left[\sum_k K_p^{\lambda(p)_k}(m_{j_1}m_{j_2})_p^{\lambda(p)_k}\right]\end{aligned}$$

We will make now two key observations, the first one is that for any subset $\mathcal{Q}_s=\{m_{j_1},m_{j_2},\ldots,m_{j_s}\}\in\mathcal{R}=\{m_1,m_2,m_3,\ldots,m_n\}$ and a fixed value of i, the i^{th} binomial expansion over $\mathcal{R}$, $\binom{m_1+\ldots+m_n}{i}$, will generate *the same* coefficient $K_p^{\lambda(p)_k}$ for the product of elements of $\mathcal{Q}_s$ corresponding to such partition, and it also does over any subset of intermediate size $s<t<n$, $\mathcal{Q}_t=\{m_{j_1},m_{j_2},\ldots,m_{j_t}\}$.

This allows a convenient simplification of the nomenclature since these coefficients will not affect the forthcoming sieve: in the former example $\mathcal{Q}_2=\{m_{j_1},m_{j_2}\}$, the sum of all products of two distinct elements arosen to all possible powers, that is the sums of products over subsets of size two, will be referred more compactly as $[2\,tuples]^{2\leqslant p\leqslant i}$ of just $2\,tuples$.

$$\textstyle\sum_{J\in\binom{1,\ldots,n}{2}}\sum_{p=2}^{i}\left[\sum_k K_p^{\lambda(p)_k}(m_{j_1}m_{j_2})_p^{\lambda(p)_k}\right]\longmapsto[2-tuples]^{2\leqslant p\leqslant i}\ \text{ or }2\,tuples$$

On the other hand, it can be observed that the sums corresponding to the subset of maximal size, that is of the size of the binomial product $s = i$, have a coefficient $K_i^{(\overbrace{1,1,...,1}^{n})} = 1$ and the $[i\,tuples]^{p=i} = \sum_{J\in\binom{1,2,\dots n}{i}} \prod_{j\in J} m_j = e_i(n)$ are exactly the *elementary symmetric polynomial* of order i.

These two facts allow naturally the following construction where coinciding typologies of $[t\,tuples]^{t\leqslant p\leqslant i}$, corresponding to $\mathcal{Q}_t \in R$, are ordered by reverse inclusion horizontally, and exploit afterwards this coincidence to extract $e_i(n)$ from $\binom{m_1+\dots+m_n}{i}$.

$$\binom{m_1+\dots+m_n}{i} \quad \ni \quad \binom{m_{j_1}+\dots+m_{j_{i-1}}}{i} \ni \binom{m_{j_1}+\dots+m_{j_{i-2}}}{i} \ni \binom{m_{j_1}+\dots+m_{j_{i-3}}}{i} \ni \cdots$$

$\mathcal{Q}_i$	$e_i(n)$			
$\mathcal{Q}_{i-1}$	$[(i-1)\ tuples]^{\,i-1\leqslant p\leqslant i}$	$[(i-1)\ tuples]^{\,i-1\leqslant p\leqslant i}$		
$\mathcal{Q}_{i-2}$	$[(i-2)\ tuples]^{\,i-2\leqslant p\leqslant i}$	$[(i-2)\ tuples]^{\,i-2\leqslant p\leqslant i}$	$[(i-2)\ tuples]^{\,i-2\leqslant p\leqslant i}$	
$\mathcal{Q}_{i-3}$	$[(i-3)\ tuples]^{\,i-3\leqslant p\leqslant i}$	$[(i-3)\ tuples]^{\,i-3\leqslant p\leqslant i}$	$[(i-3)\ tuples]^{\,i-3\leqslant p\leqslant i}$	$[(i-3)\ tuples]^{\,i-3\leqslant p\leqslant i}$
$\mathcal{Q}_4$	$[4\ tuples]^{\,4\leqslant p\leqslant i}$	$[4\ tuples]^{\,4\leqslant p\leqslant i}$	$[4\ tuples]^{\,4\leqslant p\leqslant i}$	$[4\ tuples]^{\,4\leqslant p\leqslant i}$
$\mathcal{Q}_3$	$[3\ tuples]^{\,3\leqslant p\leqslant i}$	$[3\ tuples]^{\,3\leqslant p\leqslant i}$	$[3\ tuples]^{\,3\leqslant p\leqslant i}$	$[3\ tuples]^{\,3\leqslant p\leqslant i}$
$\mathcal{Q}_2$	$[2\ tuples]^{\,2\leqslant p\leqslant i}$	$[2\ tuples]^{\,2\leqslant p\leqslant i}$	$[2\ tuples]^{\,2\leqslant p\leqslant i}$	$[2\ tuples]^{\,2\leqslant p\leqslant i}$
$\mathcal{Q}_1$	$[1\ tuples]^{\,1\leqslant p\leqslant i}$	$[1\ tuples]^{\,1\leqslant p\leqslant i}$	$[1\ tuples]^{\,1\leqslant p\leqslant i}$	$[1\ tuples]^{\,1\leqslant p\leqslant i}$

Now, in order to generate all the varieties of $s\,tuples$ of the first column in any of the others, call it $\binom{m_{j_1}+\dots+m_{j_s}}{i}$, it is necessary to add up all binomials whose numerators cover all possible sums of size s of elements within the root set $\mathcal{R} = \{m_1, m_2, m_3, \dots, m_n\}$, that is

$$\sum_{\substack{j\in J \\ J\in\binom{1,\dots,n}{s}}} \binom{\sum m_j}{i}$$

Every column has exactly the same typologies of $t\,tuples$, but not the same quantity except for $t = s$, which appear once. Indeed, observe that all $t\,tuples$ for $1 \leqslant t < s$ will be repeated as many times as sets of size $p - t$ can be formed with the n elements of the root set, from which the t that remain fixed are excluded, and in consequence every set of $t\,tuples$ appear with multiplicity $\binom{n-t}{s-t}$.

$$\binom{\overbrace{\blacktriangle+\dots+\blacktriangle}^{t}+\overbrace{\triangle+\dots+\triangle}^{s-t}}{i} \longrightarrow \binom{n-t}{s-t} posibilities; \qquad \left[\sum_{\substack{j\in J \\ J\in\binom{1,\dots,n}{s}}} \binom{\sum m_j}{i}\right]$$



Finally, all sums contain the same typologies of elements that the others and the same typologies that the expansion of $\binom{m_1+\ldots+m_n}{i}$ whose multiplicity is 1 for all t *tuples*; the elementary symmetric polynomial $e_i(n)$ can be then isolated by extracting all t *tuples* for $1 \leqslant t \leqslant i-1$ of the other columns weighted by the adequate coefficients C_h

$$e_i(n) = \binom{\sum_{i=1}^{n} m_i}{i} - \left[\sum_{h=1}^{i-1} C_h \left[\sum_{J\in\binom{1,2,\ldots n}{i-h}} \binom{\sum_{j\in J} m_j}{i}\right]\right]$$

$$e_i(n) = \binom{\sum_{j=1}^{n} m_j}{i} - \left[\left[\sum_{\substack{j\in J \\ J\in\binom{1,\ldots,n}{i-1}}}\binom{\sum m_j}{i}\right]C_1 + \left[\sum_{\substack{j\in J \\ J\in\binom{1,\ldots,n}{i-2}}}\binom{\sum m_j}{i}\right]C_2 + \left[\sum_{\substack{j\in J \\ J\in\binom{1,\ldots,n}{i-3}}}\binom{\sum m_j}{i}\right]C_3 + \left[\sum_{\substack{j\in J \\ J\in\binom{1,\ldots,n}{i-4}}}\binom{\sum m_j}{i}\right]C_4 + \left[\sum_{\substack{j\in J \\ J\in\binom{1,\ldots,n}{i-5}}}\binom{\sum m_j}{i}\right]C_5 + \cdots\right]$$



Since the typologies of the objects are the same, suffice it to consider the multiplicities to tackle the computation of the C_h and hence the following series of expressions can be constructed:

$$\begin{aligned}
1 &= \tbinom{n-i+1}{0}C_1 \\
1 &= \tbinom{n-i+2}{1}C_1 + \tbinom{n-i+2}{0}C_2 \\
1 &= \tbinom{n-i+3}{2}C_1 + \tbinom{n-i+3}{1}C_2 + \tbinom{n-i+3}{0}C_3 \\
1 &= \tbinom{n-i+4}{3}C_1 + \tbinom{n-i+4}{2}C_2 + \tbinom{n-i+4}{1}C_3 + \tbinom{n-i+4}{0}C_4 \\
&\vdots \\
1 &= \tbinom{n-i+h}{h-1}C_1 + \tbinom{n-i+h}{h-2}C_2 + \tbinom{n-i+h}{h-3}C_3 + \tbinom{n-i+h}{h-4}C_4 + \cdots + \tbinom{n-i+h}{1}C_{h-1} + \tbinom{n-i+h}{0}C_h
\end{aligned}$$

where the h^{th} term can be defined as the **recurrence** of degree $h-1$:

$$C_h = 1 - \sum_{k=1}^{h-1} C_k \binom{n-i+h}{h-k}$$

or otherwise be inserted in the **complete convolution**

$$1 = \sum_{k=1}^{h} C_k \binom{n-i+h}{h-k} \tag{2.1}$$

We sill resolve the sequence of coefficients $C_1, C_2, C_3, \ldots, C_h$ twice: first in a very immediate but scarcely elucidating way by considering a degeneration of the Vandermonde's identity and then, alternatively, in analytical terms through formal power series and generating functions.

1. Consider the complete convolution (2.1) for the term C_{h+1}

$$1 = \sum_{k=1}^{h+1} C_k \binom{n-i+h+1}{h+1-k} = \sum_{k=0}^{h} C_{k+1} \binom{n-i+h+1}{h-k}$$

and the Vandermonde's identity $\binom{l+m}{h} = \sum_{k=0}^{h} \binom{l}{k}\binom{m}{h-k}$. The first expression can be regarded as a "limit" specialization of the second one for the values $l = -n+i-1$ and $m = n-i+h+1$

$$\binom{h}{h} = 1 = \sum_{k=0}^{h} \binom{-n+i-1}{k}\binom{n-i+h+1}{h-k}$$

From where the coefficients can be pairwise identified with the numbers

$$\begin{aligned} C_k &= \binom{-n+i-1}{k-1} \\ &= (-1)^{k-1}\left(\!\!\binom{n-i+1}{k-1}\!\!\right) \end{aligned}$$

2. Let us consider again the complete convolution (2.1) for the term C_h and associate formal power series to the terms of the sequence

$$\begin{aligned} 1 &= \sum_{k=1}^{h} C_k \binom{n-i+h}{h-k} \\ \sum_{h=1}^{\infty} x^{h-1} &= \sum_{h=1}^{\infty}\left[\sum_{k=1}^{h} C_k \binom{n-i+h}{h-k}\right] x^{h-1} \\ &= \sum_{k=1}^{h} C_k \sum_{h=k}^{\infty} \binom{n-i+h}{h-k} x^{h-1} \end{aligned}$$

with the infinite geometric series on the left hand side and the binomial series on the right hand side,

$$\begin{aligned} \frac{1}{1-x} &= \sum_{k=1}^{h} C_k \frac{x^{k-1}}{(1-x)^{n-i+k+1}} \\ &= \frac{1}{x(1-x)^{n-i+1}} \sum_{k=1}^{h} C_k \left(\frac{x}{(1-x)}\right)^k \end{aligned}$$

and

$$x(1-x)^{n-i} = \sum_{k=1}^{h} C_k \left(\frac{x}{(1-x)}\right)^k$$

performing the transformation $x \longmapsto \frac{x}{(1+x)}$ the equality yields a form that admits a term by term identification with the expansion of the left hand side function

$$\begin{aligned} \frac{x}{(1+x)^{n-i+1}} &= \sum_{k=1}^{h} C_k x^k \\ \sum_{k=0}^{\infty}(-1)^k \binom{n-i+k+1}{k} x^{k+1} &= \sum_{k=0}^{\infty} C_{k+1} x^{k+1} \end{aligned}$$

from where

$$\begin{aligned} C_k &= (-1)^{k-1}\binom{n-i+k}{k-1} \\ &= \binom{-n+i-1}{k-1} \\ &= (-1)^{k-1}\left(\!\!\binom{n-i+1}{k-1}\!\!\right) \end{aligned}$$

□

References

[1] Leonhard Euler, Various analytic observations on combinations, Comentarii academiae scientiarum Petropolitanae, 13, 64-93 (1751)

[2] Henry W. Gould, The Girard-Waring power sum formulas for symmetric functions and Fibonacci sequences, Fibonacci Quarterly, 37(2), 135-140 (1999)